arXiv:math/0604016v1 [math.QA] 2 Apr 2006

MIIT-TH/01-06

The asymptotic behavior of q-exponentials and q^2-Bessel functions

V.-B.K.Rogov
MIIT, 127994, Moscow, Russia
vrogov@cemi.rssi.ru

Abstract

The connections between q^2-Bessel functions of three types and q-exponential of three types are established. The q-exponentials and the q^2-Bessel functions are represented as the Laurent series. The asymptotic behaviour of the q-exponentials and the q^2-Bessel functions is investigated.

1 Introduction

The main goal of this paper is an investigation of the asymptotic behavior of the q^2-Bessel functions for the large value of the argument. To solve this problem we establish connections between the q-exponentials and the q^2-Bessel functions.

The q-exponentials of type 1 and 2 are well-known. Although the third q-exponential is not very familiar nevertheless its properties can be received from the properties of first and second q-exponentials. The situation with the q^2-Bessel function of type 3 (the Hahn-Exton function) is similar.

Let q be the real number from $(0,1)$. The q-exponentials and the q^2-Bessel functions are the solutions of the difference equations, and so to consider them on the q-lattice (or q^2-lattice) is naturally. On the other hand these functions are determined as the convergent series in some region, and so they are analytic ones in the corresponding regions. So we will consider the continuous functions of $z > 0$ and $z = q^{n+\nu},\ n \in \mathbb{Z},\ 0 \le \nu < 1$. We will investigate the asymptotic behavior of functions $f(z)$ if $z \to \infty$ or $z = q^{n+\nu}$ and $n \to -\infty$.

The main result is contained in the Propositions 4.4, 6.1 and in the formulas (7.4) - (7.7).

We will use the standard notations

$$(a,q)_n = (1-a)(1-aq)...(1-aq^{n-1}), \quad (a,q)_\infty = \lim_{n\to\infty}(a,q)_n,$$

$$(a_1,...,a_k;q)_n = (a_1,q)_n...(a_k,q)_n,$$

and the basic hypergeometric series

$$ {}_r\Phi_s(a_1, ..., a_r; b_1, ..., b_s; q; z) = \sum_{n=0}^{\infty} \frac{(a_1, ..., a_r; q)_n}{(q, q)_n (b_1, ..., b_s; q)_n} ((-1)^n q^{\frac{n(n-1)}{2}})^{s-r+1} z^n \qquad (1.1) $$

Let $u = (1 - q^2)z$. There are three types of the q^2-exponentials:

$$ 1.\ e_q^{(1)}(u) = e_q((1 - q^2)z) = \frac{1}{((1 - q^2)z, q)_\infty} = \sum_{n=0}^{\infty} \frac{(1 - q^2)^n z^n}{(q, q)_n}, \qquad |z| < \frac{1}{1 - q^2}, \qquad (1.2) $$

$$ 2.\ e_q^{(2)}(u) = E_q((1 - q^2)z) = (-(1 - q^2)z, q)_\infty = \sum_{n=0}^{\infty} \frac{q^{\frac{n(n-1)}{2}} (1 - q^2)^n z^n}{(q, q)_n}, \qquad (1.3) $$

$$ 3.\ e_q^{(3)}(u) = {}_1\Phi_1(0; -\sqrt{q}; \sqrt{q}, -(1 - q^2)z) = \sum_{n=0}^{\infty} \frac{q^{\frac{n(n-1)}{4}} (1 - q^2)^n z^n}{(q, q)_n}. \qquad (1.4) $$

Obviously $e_q^{(1)}(u)$ is a meromorphic function and has simple poles at the points $u = q^{-n}, \ n = 0, 1, ...$. Consequently, $e_q^{(2)}(u)$ and $e_q^{(3)}(u)$ are the holomorphic functions.

Remark that for any $j = 1, \ 2, \ 3$

$$ \lim_{q \to 1-0} e_q^{(j)}((1 - q^2)z) = e^{2z}. $$

Let

$$ D_z f(z) = \frac{f(z) - f(qz)}{(1 - q^2)z}. $$

The q-exponentials satisfy the next difference equations:

$$ D_z e_q^{(1)}(u) = e_q^{(1)}(u), $$

$$ D_z e_q^{(2)}(u) = e_q^{(2)}(qu), $$

$$ D_z e_q^{(3)}(u) = e_q^{(3)}(q^{\frac{1}{2}} u). $$

The $q - \Gamma$-function is determined as

$$ \Gamma_q(\alpha) = \frac{(q, q)_\infty}{(q^\alpha, q)_\infty} (1 - q)^{1-\alpha}. $$

2 The q^2-Bessel functions

Remind that the q^2-Bessel functions of type $j, \ j = 1, 2, 3$, are determined by the following series [1]

$$ J_\nu^{(j)}(2(1 - q^2)z; q^2) = \frac{1}{\Gamma_{q^2}(\nu + 1)} \sum_{n=0}^{\infty} (-1)^n \frac{q^{(2-\delta)n(n+\nu)} (1 - q^2)^{2n} z^{\nu+2n}}{(q^2, q^2)_n (q^{2\nu+2}, q^2)_n}, $$

where the parameters j and δ are connected by formula

$$j = -\frac{3}{2}\delta^2 + \frac{5}{2}\delta + 2.$$

For $j = 1,\ |z| < \frac{1}{1-q^2}$. The modified q^2-Bessel functions are

$$I_\nu^{(j)}(2(1-q^2)z;q^2) = \frac{1}{\Gamma_{q^2}(\nu+1)} \sum_{n=0}^{\infty} \frac{q^{(2-\delta)n(n+\nu)}(1-q^2)^{2n} z^{\nu+2n}}{(q^2,q^2)_n (q^{2\nu+2},q^2)_n}. \tag{2.1}$$

Obviously

$$I_\nu^{(j)}(ze^{-i\frac{\pi}{2}};q^2) = e^{-i\frac{\nu}{2}\pi} J_\nu^{(j)}(z;q^2), \quad I_\nu^{(j)}(ze^{i\frac{\pi}{2}};q^2) = e^{i\frac{\nu}{2}\pi} J_\nu^{(j)}(z;q^2), \tag{2.2}$$

If ν is not integer the q^2-Neumann functions and the q^2-Macdonald functions respectively

$$Y_\nu^{(j)}(2(1-q^2)z;q^2) = \frac{q^{-\nu^2+\nu}}{\pi} \Gamma_{q^2}(\nu)\Gamma_{q^2}(1-\nu)[\cos\nu\pi J_\nu^{(j)}(2(1-q^2)z;q^2) - J_{-\nu}^{(j)}(2(1-q^2)z;q^2)],$$

$$K_\nu^{(j)}(2(1-q^2)z;q^2) = \frac{q^{-\nu^2+\nu}}{2} \Gamma_{q^2}(\nu)\Gamma_{q^2}(1-\nu)[I_{-\nu}^{(j)}(2(1-q^2)z;q^2) - I_\nu^{(j)}(2(1-q^2)z;q^2)].$$

q^2-Wronskian of two solutions of a difference equation of second order is determined as

$$W(f_1,f_2) = f_1(z)f_2(qz) - f_1(qz)f_2(z).$$

Proposition 2.1 *The functions* $J_\nu^{(j)}(2(1-q^2)z;q^2)$ *and* $Y_\nu^{(j)}(2(1-q^2)z;q^2)$ *satisfy the difference equation*

$$f(q^{-1}z) - (q^{-\nu}+q^\nu)f(z) + f(qz) = -q^{-\delta}(1-q^2)^2 z^2 f(q^{1-\delta}z), \tag{2.3}$$

and form the fundamental system of the solutions of this equation with q^2*-Wronskian*

$$W(J_\nu^{(j)},Y_\nu^{(j)}) = \begin{cases} \frac{q^{-\nu^2}(1-q^2)}{\pi} e_{q^2}(-(1-q^2)^2z^2) & for \quad \delta = 2 \\ \frac{q^{-\nu^2}(1-q^2)}{\pi} & for \quad \delta = 1 \\ \frac{q^{-\nu^2}(1-q^2)}{\pi} E_{q^2}((1-q^2)^2z^2) & for \quad \delta = 0. \end{cases}$$

Proof. The first statement is checked directly.

Now consider the Wronskian

$$W(J_\nu^{(j)}((1-q^2)z;q^2), Y_\nu^{(j)}((1-q^2)z;q^2)) = W(J_\nu^{(j)},Y_\nu^{(j)})(z) =$$

$$J_\nu^{(j)}((1-q^2)z;q^2)Y_\nu^{(j)}((1-q^2)qz;q^2) - J_\nu^{(j)}((1-q^2)qz;q^2)Y_\nu^{(j)}((1-q^2)z;q^2).$$

Let $\delta = 2 (j = 1)$. Because the functions $J_\nu^{(1)}$ and $Y_\nu^{(1)}$ satisfy to the equation (2.3) we have

$$J_\nu^{(1)}((1-q^2)q^2z;q^2) = -(1+(1-q^2)^2z^2)J_\nu^{(1)}((1-q^2)z;q^2) + (q^{-\nu}+q^\nu)J_\nu^{(1)}((1-q^2)qz;q^2)$$

and

$$Y_\nu^{(1)}((1-q^2)q^2z;q^2) = -(1+(1-q^2)^2z^2)Y_\nu^{(1)}((1-q^2)z;q^2) + (q^{-\nu}+q^\nu)Y_\nu^{(1)}((1-q^2)qz;q^2).$$

Hence

$$W(J_\nu^{(1)}, Y_\nu^{(1)})(qz) = (1+(1-q^2)^2z^2)W(J_\nu^{(1)}, Y_\nu^{(1)})(z),$$

i.e.

$$W(J_\nu^{(1)}, Y_\nu^{(1)})(z) = Ce_{q^2}(-(1-q^2)^2z^2).$$

By setting $z=0$, we have

$$C = \frac{q^{-\nu^2}(1-q^2)}{\pi}$$

and

$$W(J_\nu^{(1)}, Y_\nu^{(1)})(z) = \frac{q^{-\nu^2}(1-q^2)}{\pi}e_{q^2}(-(1-q^2)^2z^2).$$

The proof of the Proposition for $\delta = 1$ and 0 is similar. □

Proposition 2.2 *The functions* $I_\nu^{(j)}(2(1-q^2)z;q^2)$ *and* $K_\nu^{(j)}(2(1-q^2)z;q^2)$ *satisfy the difference equation*

$$f(q^{-1}z) - (q^{-\nu}+q^\nu)f(z) + f(qz) = q^{-\delta}(1-q^2)^2z^2f(q^{1-\delta}z),$$

and form the fundamental system of solutions of this equation with q^2*-Wronskian*

$$W(I_\nu^{(j)}, K_\nu^{(j)}) = \begin{cases} \frac{q^{-\nu^2}(1-q^2)}{2}e_{q^2}((1-q^2)^2z^2) & for \quad \delta=2 \\ \frac{q^{-\nu^2}(1-q^2)}{2} & for \quad \delta=1 \\ \frac{q^{-\nu^2}(1-q^2)}{2}E_{q^2}(-(1-q^2)^2z^2) & for \quad \delta=0. \end{cases}$$

The *Proof* is the same as above.

3 The Laurent series associated with the q-exponentials

Let z be a real positive number, and $u = e^{i\theta}z(1-q^2)$. Consider the products

$$\Lambda^{(j)}(u) = e_q^{(j)}(u)e_q^{(j)}(\frac{q}{u}); \qquad j = 1,\ 2,\ 3. \tag{3.1}$$

Proposition 3.1 *The functions* $\Lambda^{(j)}(u)$ *(3.1) can be represented as the Laurent series*

$$\Lambda^{(j)}(u) = \sum_{l=-\infty}^{\infty} u^l q^{\frac{2-\delta}{4}l^2-\frac{l}{2}} I_l^{(j)}(2q^{\frac{\delta}{4}};q). \tag{3.2}$$

If $j=1,\ q<|u|<1.$

Proof. Let $j = 1$ and $q < |u| < 1$. Substitute (1.2) into (3.1).

$$\Lambda^{(1)}(u) = \sum_{n=0}^{\infty} \frac{u^n}{(q,q)_n} \sum_{m=0}^{\infty} \frac{q^m u^{-m}}{(q,q)_m} = \sum_{l=1}^{\infty} u^{-l} \sum_{k=0}^{\infty} \frac{q^{k+l}}{(q,q)_k (q,q)_{k+l}} + \sum_{l=0}^{\infty} u^l \sum_{k=0}^{\infty} \frac{q^k}{(q,q)_k (q,q)_{k+l}} =$$

$$\sum_{l=1}^{\infty} \frac{q^l u^{-l}}{(q,q)_l} \sum_{k=0}^{\infty} \frac{q^k}{(q,q)_k (q^{l+1},q)_k} + \sum_{l=0}^{\infty} \frac{u^l}{(q,q)_l} \sum_{k=0}^{\infty} \frac{q^k}{(q,q)_k (q^{l+1},q)_k}. \tag{3.3}$$

It follows from (2.1) that if $q < |u| < 1$

$$\sum_{k=0}^{\infty} \frac{q^k}{(q,q)_k (q^{l+1},q)_k} = (q,q)_l q^{-\frac{l}{2}} I_l^{(1)}(2\sqrt{q}; q).$$

Since $I_{-l}^{(j)}(z;q) = I_l^{(j)}(z;q)$ we have

$$\Lambda^{(1)}(u) = \sum_{l=-\infty}^{\infty} u^l q^{-\frac{l}{2}} I_l^{(1)}(2\sqrt{q}; q).$$

If $j = 2$ or 3 for any $u \neq 0$ we have

$$\Lambda^{(j)}(u) = \sum_{n=0}^{\infty} \frac{q^{\frac{2-\delta}{4}n(n-1)} u^n}{(q,q)_n} \sum_{m=0}^{\infty} \frac{q^{\frac{2-\delta}{4}m(m-1)+m} u^{-m}}{(q,q)_m} =$$

$$\sum_{l=1}^{\infty} u^{-l} \sum_{k=0}^{\infty} \frac{q^{\frac{2-\delta}{4}k(k-1)+\frac{2-\delta}{4}(k+l)(k+l-1)+k+l}}{(q,q)_k (q,q)_{k+l}} + \sum_{l=0}^{\infty} u^l \sum_{k=0}^{\infty} \frac{q^{\frac{2-\delta}{4}k(k-1)+\frac{2-\delta}{4}(k+l)(k+l-1)+k}}{(q,q)_k (q,q)_{k+l}} =$$

$$\sum_{l=1}^{\infty} \frac{q^{\frac{2-\delta}{4}l(l-1)+l} u^{-l}}{(q,q)_l} \sum_{k=0}^{\infty} \frac{q^{\frac{2-\delta}{2}k(k+l)+\frac{\delta}{2}k}}{(q,q)_k (q^{l+1},q)_k} + \sum_{l=1}^{\infty} \frac{q^{\frac{2-\delta}{4}l(l-1)} u^l}{(q,q)_l} \sum_{k=0}^{\infty} \frac{q^{\frac{2-\delta}{2}k(k+l)+\frac{\delta}{2}k}}{(q,q)_k (q^{l+1},q)_k}. \tag{3.4}$$

It follows from (2.1) that for any u

$$\sum_{k=0}^{\infty} \frac{q^{\frac{2-\delta}{2}k(k+l)+k\frac{\delta}{2}}}{(q,q)_k (q^{l+1},q)_k} = (q,q)_l q^{-l\frac{\delta}{4}} I_l^{(j)}(2q^{\frac{\delta}{4}}; q).$$

So

$$\Lambda^{(j)}(u) = \sum_{l=-\infty}^{\infty} u^l q^{\frac{2-\delta}{4}l^2-\frac{l}{2}} I_l^{(j)}(2q^{\frac{\delta}{4}}; q). \tag{3.5}$$

Obviously for $j = 1$ $(\delta = 2)$ we get (3.2) from (3.5). □

Corollary 3.1 *If* u *is real and* $q < u < 1$

$$\Lambda^{(2)}(u) \leq \Lambda^{(3)}(u) \leq \Lambda^{(1)}, \tag{3.6}$$

and for $u > 1$

$$\Lambda^{(2)}(u) \leq \Lambda^{(3)}(u). \tag{3.7}$$

The *Proof* follows from the comparison of the coefficients in (3.3) and (3.4) for $j = 1,\ 2,\ 3$ $(\delta = 2,\ 0,\ 1)$. □

4 The behaviour of the q-exponentials for the large value of the argument

Proposition 4.1 *The function $\Lambda^{(1)}(u)$ (3.1) for $j=1$ satisfies the equation*

$$\Lambda^{(1)}(qu) + u\Lambda^{(1)}(u) = 0 \tag{4.1}$$

and has the form

$$\Lambda^{(1)}(u) = u^{\frac{1}{2}\log_q u - \frac{1}{2} + \frac{i\pi}{\ln q}}. \tag{4.2}$$

Proof. Consider the function $\Lambda^{(1)} = \frac{1}{(u,q)_\infty(\frac{q}{u},q)_\infty}$.

$$D_z\Lambda^{(1)}(u)e^{-i\theta} = u^{-1}\left(\Lambda^{(1)}(u) - \Lambda^{(1)}(qu)\right) = \frac{1}{u}\left(\frac{1}{(u,q)_\infty(\frac{q}{u},q)_\infty} - \frac{1}{(qu,q)_\infty(\frac{1}{u},q)_\infty}\right) =$$

$$\frac{(1-\frac{1}{u})}{u(1-u)}\frac{1}{(qu,q)_\infty(\frac{1}{u},q)_\infty} - \frac{1-u}{u(1-\frac{1}{u})}\frac{1}{(u,q)_\infty(\frac{q}{u},q)_\infty} = -u^{-2}\Lambda^{(1)}(qu) + \Lambda^{(1)}(u)$$

The last equality gives us (4.1).

The fact that (4.2) satisfies (4.1) can be checked directly. □

Proposition 4.2 *The function $\Lambda^{(2)}(u)$ (3.1) for $j=2$ satisfies the equation*

$$u\Lambda^{(2)}(qu) - \Lambda^{(2)}(u) = 0 \tag{4.3}$$

and has the form

$$\Lambda^{(2)}(u) = u^{-\frac{1}{2}\log_q u + \frac{1}{2}}. \tag{4.4}$$

Proof. Consider the function $\Lambda^{(2)} = (-u,q)_\infty(-\frac{q}{u},q)_\infty$.

$$D_z\Lambda^{(2)}(u)e^{-i\theta} = u^{-1}\left(\Lambda^{(2)}(u) - \Lambda^{(2)}(qu)\right) = \frac{1}{u}\left((-u,q)_\infty(-\frac{q}{u},q)_\infty - (-qu,q)_\infty(-\frac{1}{u},q)_\infty\right) =$$

$$\frac{1+u}{u(1+\frac{1}{u})}((-qu,q)_\infty(-\frac{1}{u},q)\infty - \frac{1+\frac{1}{u}}{u(1+u)}(-u,q)_\infty(-\frac{q}{u},q)\infty) =$$

$$\Lambda^{(2)}(qu) - u^{-2}\Lambda^{(2)}(u).$$

The last equality gives us (4.3)

The fact that (4.4) satisfies (4.3) can be checked directly. □

Represent the function $\Lambda^{(3)}(u)$ (3.1) for $j=3$ by product

$$\Lambda^{(3)}(u) = \Lambda(u)\Lambda(\frac{q}{u}) \tag{4.5}$$

and let

$$D_z\Lambda(u) = \Lambda(q^{\frac{1}{2}}u), \qquad D_z\Lambda(\frac{q}{u}) = -\frac{1}{u^2}\Lambda(\frac{q^{\frac{1}{2}}}{u}).$$

Then

$$D_z(\Lambda(u)\Lambda(\frac{q}{u})) = -\frac{1}{u^2}\Lambda(u)\Lambda(\frac{q^{\frac{1}{2}}}{u}) + \Lambda(q^{\frac{1}{2}}u)\Lambda(\frac{1}{u}).$$

Proposition 4.3 *The function* $\Lambda(u)$ *satisfies the equation*

$$\Lambda(u)\Lambda(\frac{q}{u}) - \Lambda(qu)\Lambda(\frac{1}{u}) - u\Lambda(q^{\frac{1}{2}}u)\Lambda(\frac{1}{u}) + \frac{1}{u}\Lambda(u)\Lambda(\frac{q^{\frac{1}{2}}}{u}) = 0 \tag{4.6}$$

and has the forms

$$\Lambda_1(u) = u^{\log_q u - \frac{3}{2} + \frac{2i\pi}{\ln q}} \tag{4.7}$$

or

$$\Lambda_2(u) = u^{-\frac{1}{3}\log_q u + \frac{1}{2}}. \tag{4.8}$$

Proof. Obviously

$$D_z\Lambda^{(3)}(u)e^{-i\theta} = u^{-1}\left[\Lambda(u)\Lambda(\frac{q}{u}) - \Lambda(qu)\Lambda(\frac{1}{u})\right] = \Lambda(q^{\frac{1}{2}}u)\Lambda(\frac{1}{u}) - u^{-2}\Lambda(u)\Lambda(\frac{q^{\frac{1}{2}}}{u}).$$

So we have (4.6).

The fact that (4.7) and (4.8) satisfy (4.6) can be checked directly. □

Now investigate the behaviour of the q-exponentials for the large value of the argument. Consider the cases $j = 1,\ 2,\ 3$ ($\delta = 2,\ 0,\ 1$) separately.

1. $j = 1$. Consider (4.2). Put $u = e^{i\theta}|u|,\ |u| = q^{n+\lambda},\ n = \left[\frac{\ln|u|}{\ln q}\right],\ [a]$ - is an integral part of a, and $0 \le \lambda < 1$. Then

$$\Lambda^{(1)}(u) = C^{(1)}e^{i(\theta+\pi)n}q^{\frac{1}{2}n(n-1)+\lambda n},$$

where $C^{(1)}$ is an independent on n constant. Assuming $n = 0$ we obtain

$$C^{(1)} = \Lambda^{(1)}(e^{i\theta}q^{\lambda}).$$

On the other hand it follows from (1.2) that

$$\Lambda^{(1)}(u) = \frac{1}{(u,q)_\infty(qu^{-1},q)_\infty} = \frac{1}{(q^{n+\lambda}e^{i\theta},q)_\infty(q^{1-n-\lambda}e^{-i\theta},q)_\infty} =$$

$$\frac{(q^{\lambda}e^{i\theta},q)_n}{(q^{\lambda}e^{i\theta},q)_\infty(q^{1-n-\lambda}e^{-i\theta},q)_n(q^{1-\lambda}e^{-i\theta},q)_\infty} = \frac{q^{\lambda n+\frac{1}{2}n(n-1)}e^{in\lambda(\theta+\pi)}}{(q^{\lambda}e^{i\theta},q)_\infty(q^{1-\lambda}e^{-i\theta},q)_\infty}.$$

Thus

$$C^{(1)} = \frac{1}{(q^{\lambda}e^{i\theta},q)_\infty(q^{1-\lambda}e^{-i\theta},q)_\infty}$$

and for any $u \neq 0$ [2]

$$e_q^{(1)}(u) = \frac{\Lambda^{(1)}(u)}{e_q^{(1)}(\frac{q}{u})} = \frac{q^{\lambda n)+\frac{1}{2}n(n-1)}e^{i(\pi+\theta)n}(q^{1-n-\lambda}e^{-i\theta},q)_\infty}{(q^{\lambda}e^{i\theta},q)_\infty(q^{1-\lambda}e^{-i\theta},q)_\infty}. \tag{4.9}$$

2. $j = 2$. Consider (4.4)

$$\Lambda^{(2)}(u) = u^{-\frac{1}{2}(\log_q u - 1)}$$

and assume $|u| = q^{n+\lambda}$ again. Then

$$\Lambda^{(2)}(u) = C^{(2)} e^{-i\theta n} q^{-\frac{1}{2}n(n-1)-\lambda n)}.$$

On the other hand it follows from (1.3) that

$$\Lambda^{(2)}(u) = (-u, q)_\infty (-qu^{-1}, q)_\infty = (-q^{n+\lambda} e^{i\theta}, q)_\infty (-q^{1-n-\lambda} e^{-i\theta}, q)_\infty =$$

$$\frac{(-q^{\lambda} e^{i\theta}, q)_\infty (-q^{1-n-\lambda} e^{-i\theta}, q)_n (-q^{1-\lambda} e^{-i\theta}, q)_\infty}{(-q^{\lambda} e^{i\theta}, q)_n} =$$

$$q^{-\lambda n - \frac{1}{2}n(n-1)} e^{-i\theta n} (-q^{\lambda} e^{i\theta}, q)_\infty (-q^{1-\lambda} e^{-i\theta}, q)_\infty,$$

Thus

$$C^{(2)} = (-q^{\lambda} e^{i\theta}, q)_\infty (-q^{1-\lambda} e^{-i\theta}, q)_\infty$$

and for any $u \neq 0$

$$e_q^{(2)}(u) = \frac{\Lambda^{(2)}(u)}{e_q^{(2)}(\frac{q}{u})} = q^{-\lambda n - \frac{1}{2}n(n-1)} e^{-i\theta n} \frac{(-q^{\lambda} e^{i\theta}, q)_\infty (-q^{1-\lambda} e^{-i\theta}, q)_\infty}{(-q^{1-n-\lambda} e^{-i\theta}, q)_\infty}. \tag{4.10}$$

3. $j = 3$. The equation (4.6) has two solutions (4.7) and (4.8). Substituting these functions into (4.5) we obtain

$$\Lambda_1^{(3)}(u) = C_1 u^{2(\log_q u - 1)}, \qquad \Lambda_2^{(3)}(u) = C_2 u^{-\frac{2}{3}(\log_q u - 1)}.$$

For an arbitrary real $u > 1$ take the solution $\Lambda_2^{(3)}(u)$ and require that

$$u^{-\frac{1}{2}(\log_q u - 1)} \leq q^{\mu} u^{-\frac{2}{3}(\log_q u - 1)},$$

according to (3.7).

$$\mu - \frac{2}{3}(\log_q u - 1)\log_q u \leq -\frac{1}{2}(\log_q u - 1)\log_q u,$$

$$\mu \leq \frac{1}{6}(\log_q u - 1)\log_q u,$$

and we put $\mu = -\frac{1}{24}$. Then (3.6) is fulfilled for $u:\ q < u < 1$. So for an arbitrary complex $u \neq 0$

$$\Lambda^{(3)}(u) = q^{-\frac{1}{24}} u^{-\frac{2}{3}(\log_q u - 1)},$$

and we assume $|u| = q^{n+\lambda}$ again.

$$\Lambda^{(3)}(u) = C^{(3)} q^{-\frac{2}{3}n(n-1) - \frac{4}{3}n\lambda} e^{-\frac{4i\theta}{3}n},$$

$$C^{(3)} = q^{-\frac{1}{24}} e_q^{(3)}(q^{\{\lambda\}} e^{i\theta}) e_q^{(3)}(q^{1-\{\lambda\}} e^{-i\theta}).$$

Hence for any $u \neq 0$

$$e_q^{(3)}(u) = \frac{\Lambda^{(3)}(u)}{e_q^{(3)}(\frac{q}{u})} = q^{-\frac{2}{3}n(n-1) - \frac{4}{3}n\lambda - \frac{1}{24}} e^{-\frac{4i\theta}{3}n} \frac{e_q^{(3)}(q^{\lambda} e^{i\theta}) e_q^{(3)}(q^{1-\lambda} e^{-i\theta})}{e_q^{(3)}(q^{1-n-\lambda} e^{-i\theta})}. \tag{4.11}$$

Proposition 4.4 *The asymptotic behaviour of the q-exponentials has the form*

$$e_q^{(j)}(u) = \begin{cases} q^{\frac{1}{2}N} e^{i(\theta+\pi)n} C^{(1)}(1+o(q^{-n})) & \text{for} \quad j=1 \\ q^{-\frac{1}{2}N} e^{-i\theta n} C^{(2)}(1+o(q^{-n})) & \text{for} \quad j=2 \\ q^{-\frac{2}{3}N-\frac{1}{24}} e^{-\frac{4i\theta}{3}n} C^{(3)}(1+o(q^{-n})) & \text{for} \quad j=3, \end{cases}$$

where

$$N = n(n-1) + 2\lambda n, \tag{4.12}$$

and

$$C^{(j)} = e_q^{(j)}(q^{\lambda} e^{i\theta}) e_q^{(j)}(q^{1-\lambda} e^{-i\theta}).$$

Proof. Obviously

$$\lim_{n\to-\infty} e_q^{(j)}(q^{1-n-\lambda} e^{-i\theta}) = 1.$$

Now the statement of the Proposition follows from (4.9) - (4.11). □

5 The representations of the q^2-Bessel functions as the Laurent series

Let u be real.

In [3] it has been received the following representations for the modified q^2-Bessel functions of kind 1 and 2

$$I_\nu^{(j)}(2u;q^2) = \frac{a_\nu}{\sqrt{2u}} \left[e_q^{(j)}(u)\Phi_\nu(u) + ie^{i\nu\pi} e_q^{(j)}(-u)\Phi_\nu(-u) \right], \tag{5.1}$$

$$K_\nu^{(j)}(2u;q^2) = \frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{2a_\nu\sqrt{2u}} e_q^{(j)}(-u)\Phi_\nu(-u), \quad \text{for } \nu \neq n, \tag{5.2}$$

where

$$u = (1-q^2)z, \qquad a_\nu^2 = a_{-\nu}^2 = \begin{cases} \frac{q^{-\nu+\frac{1}{2}}(1-q^2)}{2\Gamma_{q^2}(\nu)\Gamma_{q^2}(1-\nu)\sin\nu\pi} & \text{for} \quad \nu \neq n \\ \frac{q^{-n^2+\frac{1}{2}}\ln q^{-2}}{2\pi} & \text{for} \quad \nu = n \end{cases},$$

$$\Phi_\nu(u) = {}_2\Phi_1(q^{\nu+\frac{1}{2}}, q^{-\nu+\frac{1}{2}}; -q; q, \frac{q}{u}). \tag{5.3}$$

In the case $\nu = n$ $K_n^{(j)}$ can be received as the limit of $K_\nu^{(j)}$ if ν tends to n.

It follows from (1.2), (1.3) and (5.3) that $\sqrt{u} I_\nu^{(1)}(2u;q^2)$ is a meromorphic function and has simple poles at the points $u = \pm q^{-n}$, $n = 0, 1, \ldots$. $\sqrt{u} K_\nu^{(1)}(2u;q^2)$ is a meromorphic function and has simple poles at the points $u = -q^{-n}$, $n = 0, 1, \ldots$. $\sqrt{u} I_\nu^{(2)}(2u;q^2)$ and $\sqrt{u} K_\nu^{(2)}(2u;q^2)$ are the holomorphic functions outside a neighborhood of zero (see [3])

Using (2.2) we can write

$$J_\nu^{(j)}(2u;q^2) = \frac{a_\nu}{\sqrt{2u}} \left[e^{-i(\frac{\pi}{4}+\nu\frac{\pi}{2})} e_q^{(j)}(iu)\Phi_\nu(iu) + ie^{i(\frac{\pi}{4}+\nu\frac{\pi}{2})} e_q^{(j)}(-iu)\Phi_\nu(-iu) \right], \tag{5.4}$$

$$Y_\nu^{(j)}(2u;q^2) = \tag{5.5}$$

$$-\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{\pi\sqrt{2u}a_\nu}\left[e^{i(\frac{\pi}{4}-\nu\frac{\pi}{2})}e_q^{(j)}(iu)\Phi_\nu(iu)+ie^{-i(\frac{\pi}{4}-\nu\frac{\pi}{2})}e_q^{(j)}(-iu)\Phi_\nu(-iu)\right],$$

if ν is not integer. In the case $\nu = n$ $\;Y_n^{(j)}$ can be received as the limit of $Y_\nu^{(j)}$ if ν tends to n.

The functions $\sqrt{u}J_\nu^{(1)}(2u;q^2)$ and $\sqrt{u}Y_\nu^{(1)}(2u;q^2)$ are the meromorphic functions and have simple poles at the points $u = \pm iq^{-n}$, $\;n = 0, 1, \ldots.$ The functions $\sqrt{u}J_\nu^{(2)}(2u;q^2)$ and $\sqrt{u}Y_\nu^{(2)}(2u;q^2)$ are the holomorphic functions outside a neighborhood of zero.

To receive the similar formulas for q^2-Bessel functions of kind 3 we use the following

Lemma 5.1 *Let the functions*

$$f^{(j)}(z) = \sum_{k=-\infty}^{\infty} b_k^{(j)} z^{k-\frac{1}{2}} \tag{5.6}$$

satisfy the equations

$$q^{\frac{1}{2}}f^{(j)}(q^{-1}z) - (q^{-\nu}+q^{\nu})f^{(j)}(z) + q^{-\frac{1}{2}}f^{(j)}(qz) = q^{-\frac{\delta+1}{2}}z^2 f^{(j)}(q^{1-\delta}z) \tag{5.7}$$

where $j = -\frac{3}{2}\delta^2 + \frac{5}{2}\delta + 2$, $\quad\delta = 0, 1$ *and* 2. *Then*

$$b_k^{(3)} = \sqrt{b_k^{(1)}b_k^{(2)}} \tag{5.8}$$

for any k

Proof. It follows from (5.6) and (5.7) that

$$b_k^{(j)} = \frac{b_{k-2}^{(j)}q^{k\delta-3(1-\frac{\delta}{2})}}{(1-q^{-\nu+k-\frac{1}{2}})(1-q^{\nu+k-\frac{1}{2}})} = \begin{cases} \frac{b_0^{(j)}q^{l(l\delta-3+\frac{5}{2}\delta)}}{(q^{-\nu+\frac{3}{2}},q^2)_l(q^{\nu+\frac{3}{2}},q^2)_l} & \text{for} \quad k = 2l \\ \frac{b_1^{(j)}q^{l(l\delta-3+2\delta)}}{(q^{-\nu+\frac{3}{2}},q^2)_l(q^{\nu+\frac{3}{2}},q^2)_l} & \text{for} \quad k = 2l+1, \end{cases}$$

i.e. (5.8) is fulfilled. □

Find the coefficient of function

$$\frac{1}{\sqrt{u}}e_q^{(j)}(u)\Phi_\nu(u) = \frac{1}{\sqrt{u}}\left[\sum_{l=1}^{\infty}c_{l-}^{(j)}u^{-l} + \sum_{l=0}^{\infty}c_{l+}^{(j)}u\right], \qquad j = 1, 2,$$

in their decompositions in the Laurent series. Because these functions satisfy (5.7) for $\delta = 2$ and 0 respectively the following series satisfies (5.7) for $\delta = 1$

$$\left[\sum_{l=1}^{\infty}\sqrt{c_{l-}^{(1)}c_{l-}^{(2)}}u^{-l-\frac{1}{2}} + \sum_{l=0}^{\infty}\sqrt{c_{l+}^{(1)}c_{l+}^{(2)}}u^{l-\frac{1}{2}}\right]. \tag{5.9}$$

Using (1.2), (1.3) and (5.3), we can write for $j = 1,\ 2$

$$c_{l-}^{(j)} = \sum_{k=0}^{\infty}\frac{q^{\frac{2-\delta}{4}k(k-1)}}{(q,q)_k}\frac{(q^{-\nu+\frac{1}{2}},q)_{k+l}(q^{\nu+\frac{1}{2}},q)_{k+l}}{(q^2,q^2)_{k+l}}q^{k+l} =$$

$$\frac{(q^{-\nu+\frac{1}{2}},q)_l(q^{\nu+\frac{1}{2}},q)_l}{(q^2,q^2)_l}q^l\sum_{k=0}^{\infty}\frac{(q^{-\nu+l+\frac{1}{2}},q)_k(q^{\nu+l+\frac{1}{2}},q)_k}{(q^{2l+2},q^2)_k(q,q)_k}q^{\frac{2-\delta}{4}k(k-1)+k}.$$

$$c_{l+}^{(j)}=\sum_{k=0}^{\infty}\frac{q^{\frac{2-\delta}{4}(k+l)(k+l-1)}}{(q,q)_{k+l}}\frac{(q^{-\nu+\frac{1}{2}},q)_k(q^{\nu+\frac{1}{2}},q)_k}{(q^2,q^2)_k}q^k=$$

$$\frac{q^{\frac{l(l-1)}{2}}}{(q,q)_l}\sum_{k=0}^{\infty}\frac{(q^{-\nu+\frac{1}{2}},q)_k(q^{\nu+\frac{1}{2}},q)_k}{(q^2,q^2)_k(q^{l+1},q)_k}q^{\frac{2-\delta}{4}k(k+l-1)+k}.$$

For this reason we have

$$\sqrt{c_{l-}^{(1)}c_{l-}^{(2)}}=\frac{(q^{-\nu+\frac{1}{2}},q)_l(q^{\nu+\frac{1}{2}},q)_l}{(q^2,q^2)_l}q^l\times \tag{5.10}$$

$$\times\sqrt{\sum_{k=0}^{\infty}\frac{(q^{-\nu+l+\frac{1}{2}},q)_k(q^{\nu+l+\frac{1}{2}},q)_k}{(q^{2l+2},q^2)_k(q,q)_k}q^k\sum_{m=0}^{\infty}\frac{(q^{-\nu+l+\frac{1}{2}},q)_m(q^{\nu+l+\frac{1}{2}},q)_m}{(q^{2l+2},q^2)_m(q,q)_m}q^{\frac{m(m+1)}{2}}},$$

$$\sqrt{c_{l+}^{(1)}c_{l+}^{(2)}}= \tag{5.11}$$

$$\frac{q^{\frac{l(l-1)}{4}}}{(q,q)_l}\sqrt{\sum_{k=0}^{\infty}\frac{(q^{-\nu+\frac{1}{2}},q)_k(q^{\nu+\frac{1}{2}},q)_k}{(q^2,q^2)_k(q^{l+1},q)_k}q^k\sum_{m=0}^{\infty}\frac{(q^{-\nu+\frac{1}{2}},q)_m(q^{\nu+\frac{1}{2}},q)_m}{(q^2,q^2)_m(q^{l+1},q)_m}q^{\frac{m(m+l+1)}{2}}}.$$

The representations of $I_\nu^{(j)}$, $j=1,\ 2$, have the form

$$I_\nu^{(j)}(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}\times$$

$$\times\left[\sum_{l=1}^{\infty}c_{l-}^{(j)}u^{-l}+\sum_{l=0}^{\infty}c_{l+}^{(j)}u^l+ie^{i\nu\pi}\left(\sum_{l=1}^{\infty}(-1)^lc_{l-}^{(j)}u^{-l}+\sum_{l=0}^{\infty}(-1)^lc_{l+}^{(j)}u^l\right)\right]=$$

$$\frac{a_\nu}{\sqrt{2u}}\left[\sum_{l=1}^{\infty}(1+ie^{i(\nu+l)\pi})c_{l-}^{(j)}u^{-l}+\sum_{l=0}^{\infty}(1+ie^{i(\nu+l)\pi})c_{l+}^{(j)}u^l\right].$$

Hence

$$I_\nu^{(3)}(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}\times \tag{5.12}$$

$$\times\left[\sum_{l=1}^{\infty}(1+ie^{i(\nu+l)\pi})\sqrt{c_{l-}^{(1)}c_{l-}^{(2)}}u^{-l}+\sum_{l=0}^{\infty}(1+ie^{i(\nu+l)\pi})\sqrt{c_{l+}^{(1)}c_{l+}^{(2)}}u^l\right].$$

In the similar way we obtain

$$K_\nu^{(3)}(2u;q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{2a_\nu\sqrt{2u}}\left[\sum_{l=1}^{\infty}(-1)^l\sqrt{c_{l-}^{(1)}c_{l-}^{(2)}}u^{-l}+\sum_{l=0}^{\infty}(-1)^l\sqrt{c_{l+}^{(1)}c_{l+}^{(2)}}u^l\right], \tag{5.13}$$

$$J_\nu^{(3)}(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}e^{-i\frac{\pi}{4}-i\frac{\nu}{2}\pi}\times \tag{5.14}$$

$$\times\left[\sum_{l=1}^{\infty}(1+ie^{i\nu\pi})\sqrt{c_{l-}^{(1)}c_{l-}^{(2)}}u^{-l}+\sum_{l=0}^{\infty}(1+ie^{i\nu\pi})\sqrt{c_{l+}^{(1)}c_{l+}^{(2)}}u^{l}\right],$$

$$Y_\nu^{(3)}(2u;q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{a_\nu\pi\sqrt{2u}}e^{i\frac{\pi}{4}-i\frac{\nu}{2}\pi}\times \tag{5.15}$$

$$\times\left[\sum_{l=1}^{\infty}(-1+ie^{i\nu\pi})\sqrt{c_{l-}^{(1)}c_{l-}^{(2)}}u^{-l}+\sum_{l=0}^{\infty}(-1+ie^{i\nu\pi})\sqrt{c_{l+}^{(1)}c_{l+}^{(2)}}u^{l}\right].$$

6 The asymptotic behaviour of the q^2-Bessel functions of type 1 and 2

Let $u=q^{n+\lambda}$, $n=\left[\frac{\ln u}{\ln q}\right]$, where $[a]$ is an integral part of a and $0\le\lambda<0$. Introduce the designations

$$Q_{\pm}^{(j)}=\frac{e_q^{(j)}(\pm q^{\lambda})e_q^{(j)}(\pm q^{1-\lambda})}{e_q^{(j)}(\pm q^{1-n-\lambda})},\quad Q_{\pm i}^{(j)}=\frac{e_q^{(j)}(\pm iq^{\lambda})e_q^{(j)}(\mp iq^{1-\lambda})}{e_q^{(j)}(\mp iq^{1-n-\lambda})} \tag{6.1}$$

and define N by (4.12) as previously. Then it follows from (4.9) - (4.11) that

$$e_q^{(1)}(\pm u)=q^{\frac{1}{2}N}e^{i\frac{\pi}{2}(1\pm1)n}Q_{\pm}^{(1)},\qquad e_q^{(1)}(\pm iu)=q^{\frac{1}{2}N}e^{i\pi(1\pm\frac{1}{2})n}Q_{\pm i}^{(1)}.$$

$$e_q^{(2)}(\pm u)=q^{-\frac{1}{2}N}e^{i\frac{\pi}{2}(1\mp1)n}Q_{\pm}^{(2)},\qquad e_q^{(2)}(\pm iu)=q^{-\frac{1}{2}N}e^{\mp i\frac{\pi}{2}n}Q_{\pm i}^{(2)}.$$

Using (5.1) and (5.2) we obtain

$$I_\nu^{(1)}(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}q^{\frac{1}{2}N}\left[e^{in\pi}Q_{+}^{(1)}\Phi_\nu(u)+ie^{i\nu\pi}Q_{-}^{(1)}\Phi_\nu(-u)\right], \tag{6.2}$$

$$K_\nu^{(1)}(2u;q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{2a_\nu\sqrt{2u}}q^{\frac{1}{2}N}Q_{-}^{(1)}\Phi_\nu(-u), \tag{6.3}$$

$$I_\nu^{(2)}(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}q^{-\frac{1}{2}N}\left[Q_{+}^{(2)}\Phi_\nu(u)+ie^{i\nu\pi+in)\pi}Q_{-}^{(2)}\Phi_\nu(-u)\right], \tag{6.4}$$

$$K_\nu^{(2)}(2u);q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{2a_\nu\sqrt{2u}}q^{-\frac{1}{2}N}e^{i\pi n}Q_{-}^{(2)}\Phi_\nu(-u). \tag{6.5}$$

Us the formulas (5.4), (5.5) and (6.1). Then

$$J_\nu^{(1)}(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}q^{\frac{1}{2}N}\times \tag{6.6}$$

$$\times\left[e^{-i(2\nu+1)\frac{\pi}{4}+i\frac{3\pi}{2}n}Q_{+i}^{(1)}\Phi_\nu(iu)+e^{i(2\nu+1)\frac{\pi}{4}+i\frac{\pi}{2}n}Q_{-i}^{(1)}\Phi_\nu(-iu)\right]$$

$$Y_\nu^{(1)}(2u;q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{\pi a_\nu\sqrt{2u}}q^{\frac{1}{2}N}\times \tag{6.7}$$

$$\times\left[-e^{-i(2\nu-1)\frac{\pi}{4}+i\frac{3\pi}{2}n}Q^{(1)}_{+i}\Phi_\nu(iu)-e^{i(2\nu-1)\frac{\pi}{4}+i\frac{\pi}{2}n}Q^{(1)}_{-i}\Phi_\nu(-iu)\right]$$

$$J^{(2)}_\nu(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}q^{-\frac{1}{2}N}\times \tag{6.8}$$

$$\times\left[e^{-i(2\nu+1)\frac{\pi}{4}-i\frac{\pi}{2}n}Q^{(2)}_{+i}\Phi_\nu(iu)+e^{i(2\nu+1)\frac{\pi}{4}+i\frac{\pi}{2}n}Q^{(2)}_{-i}\Phi_\nu(-iu)\right]$$

$$Y^{(2)}_\nu(2u;q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{\pi a_\nu\sqrt{2u}}q^{-\frac{1}{2}N}\times \tag{6.9}$$

$$\times\left[-e^{-i(2\nu-1)\frac{\pi}{4}-i\frac{\pi}{2}n}Q^{(2)}_{+i}\Phi_\nu(iu)-e^{i(2\nu-1)\frac{\pi}{4}+i\frac{\pi}{2}n}Q^{(2)}_{-i}\Phi_\nu(-iu)\right].$$

Now we can describe the asymptotic behaviour of q^2-Bessel functions of type 1 and 2. Note that if $n\to-\infty$ $(|u|\to\infty)$,

$$\Phi_\nu(\pm u)Q^{(j)}_\pm\to e^{(j)}_q(\pm q^\lambda)e^{(j)}_q(\pm q^{1-\lambda})$$

and

$$\Phi_\nu(\pm iu)Q^{(j)}_{\pm i}\to\frac{1}{e_q}^{(j)}(\pm iq^\lambda)e^{(j)}_q(\mp iq^{1-\lambda}).$$

Proposition 6.1 *The following asymptotic formulas take place for* $I^{(j)}_\nu(2u;q^2)$, $K^{(j)}_\nu(2u;q^2)$, $J^{(1)}_\nu(2u;q^2)$, $Y^{(1)}_\nu(2u;q^2)$, $j=1,\ 2$

$$I^{(1)}_\nu(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}q^{\frac{1}{2}N}\times$$

$$\times\left[[e^{in\pi}e^{(1)}_q(q^\lambda)e^{(1)}_q(q^{1-\lambda})+ie^{i\nu\pi}e^{(1)}_q(-q^\lambda)e^{(1)}_q(-q^{1-\lambda})\right](1+o(q^{-n})),$$

$$K^{(1)}_\nu(2u;q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{2a_\nu\sqrt{2u}}q^{\frac{1}{2}N}e^{(1)}_q(-q^\lambda)e^{(1)}_q(-q^{1-\lambda})(1+o(q^{-n})),$$

$$I^{(2)}_\nu(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}q^{-\frac{1}{2}N}\left[e^{(2)}_q(q^\lambda)e^{(2)}_q(q^{1-\lambda})+\right.$$

$$\left.ie^{i\nu\pi+i\pi n}e^{(2)}_q(-q^\lambda)e^{(2)}_q(-q^{1-\lambda})\right](1+o(q^{-n})),$$

$$K^{(2)}_\nu(2u;q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{2a_\nu\sqrt{2u}}q^{-\frac{1}{2}N}e^{i\pi n}e^{(2)}_q(-q^\lambda)e^{(2)}_q(-q^{1-\lambda})(1+o(q^{-n})),$$

$$J^{(1)}_\nu(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}q^{\frac{1}{2}N}\left[e^{-i(2\nu+1)\frac{\pi}{4}+i\frac{3\pi}{2}n}e^{(1)}_q(iq^\lambda)e^{(1)}_q(-iq^{1-\lambda})+\right.$$

$$\left.e^{i(2\nu+1)\frac{\pi}{4}+i\frac{\pi}{2}(n+\lambda)}e^{(1)}_q(-iq^\lambda)e^{(1)}_q(iq^{1-\lambda})\right](1+o(q^{-n})),$$

$$Y^{(1)}_\nu(2u;q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{\pi\sqrt{2u}}q^{\frac{1}{2}N}\left[-e^{-i(2\nu-1)\frac{\pi}{4}+i\frac{3\pi}{2}n}e^{(1)}_q(iq^\lambda)e^{(1)}_q(-iq^{1-\lambda})-\right.$$

$$\left.e^{i(2\nu-1)\frac{\pi}{4}+i\frac{\pi}{2}n}e^{(1)}_q(-iq^\lambda e^{(1)}_q(iq^{1-\lambda})\right](1+o(q^{-n})),$$

$$J^{(2)}_\nu(2uz;q^2)=\frac{a_\nu}{\sqrt{2u}}q^{-\frac{1}{2}N}\left[e^{-i(2\nu+1)\frac{\pi}{4}-i\frac{\pi}{2}n}e^{(2)}_q(iq^\lambda)e^{(2)}_q(-iq^{1-\lambda})+\right.$$

$$e^{i(2\nu+1)\frac{\pi}{4}+i\frac{\pi}{2}n}e_q^{(2)}(-iq^{\lambda})e_q^{(2)}(iq^{1-\lambda})\big]\left(1+o(q^{-n})\right),$$

$$Y_\nu^{(2)}(2u;q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{\pi a_\nu\sqrt{2u}}q^{-\frac{1}{2}N}\big[-e^{-i(2\nu-1)\frac{\pi}{4}-i\frac{\pi}{2}n}e_q^{(2)}(iq^{\lambda})e_q^{(2)}(-iq^{1-\lambda})+$$

$$e^{i(2\nu-1)\frac{\pi}{4}+i\frac{\pi}{2}n}e_q^{(2)}(-iq^{\lambda})e_q^{(2)}(iq^{1-\lambda})\big]\left(1+o(q^{-n})\right).$$

Proof. The statement of Proposition follows from (6.2) - (6.9). □

7 The asymptotic behaviour of the q^2-Bessel functions of type 3

Consider the coefficient (5.10). Obviously

$$q^{-\nu+\frac{1}{2}}+q^{\nu-\frac{1}{2}}\geq 2 \quad\Rightarrow\quad q^{-\nu+\frac{1}{2}}-1\geq 1-q^{\nu-\frac{1}{2}} \quad\Rightarrow$$

$$q^{-\frac{1}{2}}(q^{-\nu+\frac{1}{2}}-1)\geq q^{\frac{1}{2}}(1-q^{\nu-\frac{1}{2}}) \quad\Rightarrow q^{-\nu}+q^{\nu}\geq q^{\frac{1}{2}}+q^{-\frac{1}{2}} \quad\Rightarrow$$

$$(q^{\nu}+q^{-\nu})q^{l+k+\frac{1}{2}}\geq (q^{\frac{1}{2}}+q^{-\frac{1}{2}})q^{2l+2k+\frac{1}{2}} \quad\Rightarrow$$

$$1-(q^{\nu}+q^{-\nu})q^{l+k+\frac{1}{2}}+q^{2l+2k+1}\leq 1-q^{2l+2k} \quad\Rightarrow$$

$$\frac{(1-q^{-\nu+l+k+\frac{1}{2}})(1-q^{\nu+l+k+\frac{1}{2}})}{1-q^{2l+2k}}\leq 1.$$

Hence it follows from (1.2) and (1.3) that

$$\sum_{k=0}^{\infty}\frac{(q^{-\nu+l+k+\frac{1}{2}},q)_k(q^{\nu+l+k+\frac{1}{2}},q)_k}{(q^{2l+2},q^2)_k(q,q)_k}q^k\leq\sum_{k=0}^{\infty}\frac{q^k}{(q,q)_k}=e_q(q),$$

$$\sum_{m=0}^{\infty}\frac{(q^{-\nu+l+m+\frac{1}{2}},q)_m(q^{\nu+l+m+\frac{1}{2}},q)_m}{(q^{2l+2},q^2)_m(q,q)_m}q^{\frac{m(m+1)}{2}}\leq\sum_{m=0}^{\infty}\frac{q^{\frac{m(m+1)}{2}}}{(q,q)_m}=E_q(q)$$

and

$$\sqrt{c_{l-}^{(1)}c_{l-}^{(2)}}\leq\sqrt{e_q(q)E_q(q)}\frac{(q^{-\nu+\frac{1}{2}},q)_l(q^{\nu+\frac{1}{2}},q)_l}{(q^2,q^2)_l}q^l.$$

But this coefficient is not interesting for us because the first summand of (5.9) tends to zero if $|u|\rightarrow\infty$.

Consider coefficient (5.11). It follows from (1.1) and (5.3) that for arbitrary $l\geq 0$ and $i=1,2,...,k$

$$1<\frac{1}{1-q^{l+i}}\leq\frac{1}{1-q},$$

and for arbitrary $i=1,2,...,m$

$$0<\frac{q^{l+1}}{1-q^{l+i}}<\frac{q}{1-q}.$$

So

$$\sum_{k=0}^{\infty}\frac{(q^{-\nu+\frac{1}{2}},q)_k(q^{\nu+\frac{1}{2}},q)_k}{(q^2,q^2)_k(q^{l+1},q)_k}q^k=\sum_{k=0}^{\infty}\frac{(q^{-\nu+\frac{1}{2}},q)_k(q^{\nu+\frac{1}{2}},q)_k}{(q^2,q^2)_k}(\alpha q)^k=$$

$${}_2\Phi_1(q^{\nu+\frac{1}{2}},q^{-\nu+\frac{1}{2}};-q;q,\alpha q)=\varphi_1^2(\alpha) \tag{7.1}$$

for some $\alpha\in(1,\frac{1}{1-q})$, and

$$\sum_{m=0}^{\infty}\frac{(q^{-\nu+\frac{1}{2}},q)_m(q^{\nu+\frac{1}{2}},q)_m}{(q^2,q^2)_m(q^{l+1},q)_m}q^{\frac{1}{2}m(m-1)+\beta m+m}=\sum_{m=0}^{\infty}\frac{(q^{-\nu+\frac{1}{2}},q)_m(q^{\nu+\frac{1}{2}},q)_m}{(q^2,q^2)_m}q^{\frac{1}{2}m(m-1)}(\beta q)^m=$$

$${}_2\Phi_2(q^{\nu+\frac{1}{2}},q^{-\nu+\frac{1}{2}};-q,0;q,-\beta q)=\varphi_2^2(\beta) \tag{7.2}$$

for some $\beta\in(0,\frac{1}{1-q})$.

Hence

$$\sqrt{c_{l+}^{(1)}c_{l+}^{(2)}}=\frac{q^{\frac{l(l-1)}{4}}}{(q,q)_l}\varphi_l(\alpha)\varphi_2(\beta)=\frac{q^{\frac{l(l-1)}{4}}}{(q,q)_l}\varphi(\alpha,\beta). \tag{7.3}$$

Now substituting (7.3) into (5.12) - (5.15) and using (1.4) we can write

$$I_\nu^{(3)}(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}\left[\sum_{l=1}^{\infty}(1+ie^{i(\nu+l)\pi})\sqrt{c_{l-}^{(1)}c_{l-}^{(2)}}u^{-l}+\sum_{l=0}^{\infty}(1+ie^{i(\nu+l)\pi})\sqrt{c_{l+}^{(1)}c_{l+}^{(2)}}u^{l}\right]=$$

$$\frac{a_\nu}{\sqrt{2u}}\sum_{l=0}^{\infty}(1+ie^{i(\nu+l)\pi})\sqrt{c_{l+}^{(1)}c_{l+}^{(2)}}u^{l}(1+o(q^{-n}))=$$

$$\frac{a_\nu}{\sqrt{2u}}\varphi(\alpha,\beta)(e_q^{(3)}(u)+ie^{i\nu\pi}e_q^{(3)}(-u))(1+o(q^{-n}))=$$

$$\frac{a_\nu}{\sqrt{2u}}q^{-\frac{2}{3}N-\frac{1}{24}}\varphi(\alpha,\beta)\left[\frac{e_q^{(3)}(q^\lambda)e_q^{(3)}(q^{1-\lambda})}{e_q^{(3)}(q^{1-n-\lambda})}+ie^{i\nu\pi}\frac{e_q^{(3)}(-q^\lambda)e_q^{(3)}(-q^{1-\lambda})}{e_q^{(3)}(-q^{1-n-\lambda})}\right](1+o(q^{-n}))$$

Because $e_q^{(3)}(-q^{1-n-\lambda})\to 1$ if $n\to-\infty$ we have

$$I_\nu^{(3)}(2u;q^2)=\frac{a_\nu}{\sqrt{2u}}q^{-\frac{2}{3}N-\frac{1}{24}}\varphi(\alpha,\beta)\times \tag{7.4}$$

$$\times\left[e_q^{(3)}(q^\lambda)e_q^{(3)}(q^{1-\lambda})+ie^{i\nu\pi}e_q^{(3)}(-q^\lambda)e_q^{(3)}(-q^{1-\lambda})\right](1+o(q^{-n}))$$

In the similar way we obtain

$$K_\nu^{(3)}(2u;q^2)=\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{2a_\nu\sqrt{2u}}q^{-\frac{2}{3}N-\frac{1}{24}}\varphi(\alpha,\beta)e_q^{(3)}(-q^\lambda)e_q^{(3)}(-q^{1-\lambda})(1+o(q^{-n})), \tag{7.5}$$

$$J_\nu^{(3)}(2u;q^2)=\frac{2a_\nu}{\sqrt{2z}}q^{-\frac{2}{3}N-\frac{1}{24}}\varphi(\alpha,\beta)e^{-i(2\nu+1)\frac{\pi}{4}}\times \tag{7.6}$$

$$\times\left[-e_q^{(3)}(iq^\lambda)e_q^{(3)}(-iq^{1-\lambda})+ie^{i\nu\pi}e_q^{(3)}(-iq^\lambda)e_q^{(3)}(iq^{1-\lambda})\right](1+o(q^{-n})),$$

$$Y_\nu^{(3)}(2u;q^2)=-\frac{q^{-\nu^2+\frac{1}{2}}(1-q^2)}{a_\nu\pi\sqrt{2u}}q^{-\frac{2}{3}N-\frac{1}{24}}e^{-i(2\nu-1)\frac{\pi}{4}}\varphi(\alpha,\beta)\times \tag{7.7}$$

$$\times\left[e_q^{(3)}(iq^\lambda)e_q^{(3)}(-iq^{1-\lambda})+ie^{i\nu\pi}e_q^{(3)}(-iq^\lambda)e_q^{(3)}(iq^{1-\lambda})\right](1+o(q^{-n})),$$

where $\varphi(\alpha,\beta)=\varphi_1(\alpha)\varphi_2(\beta)$ is determined be (7.1) and (7.2).

References

[1] G.Gasper and M.Rahman. *Basic Hypergeometric Series.* Cambridge University Press, 1990

[2] Mourad E.H.Ismail. *Some Properties of Jackson's Third q-Bessel Function.* May 28, 1996

[3] M.A.Olshanetsky and V.-B.K.Rogov *Modified q-Bessel functions and q-Macdonald functions*, Sbornik: Mathematics **187**: 10 (1996), 1525-1544.